

APPLICATION  
FOR  
UNITED STATES LETTERS PATENT

TITLE: THYMOSIN BETA 4 PROMOTES WOUND REPAIR

APPLICANT: Hynda K. Kleinman, Allan Goldstein,  
Katherine M. Malinda and Gabriel Sosne

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208

**THYMOSIN  $\beta$ 4 PROMOTES WOUND REPAIR**

**STATEMENT AS TO FEDERALLY SPONSORED RESEARCH**

This invention was made in part with funds from the National Institutes of  
5 Health, Intramural Program. The government may have certain rights in this invention.

**CROSS REFERENCE TO RELATED APPLICATIONS**

This application claims priority from Provisional Application Serial No.  
60/094,690, filed July 30, 1998, which is incorporated herein by reference in its  
entirety and to which application a priority claim is made under 35 U.S.C. §119(e).

**TECHNICAL FIELD OF THE INVENTION**

The present invention relates generally to tissue repair and more  
specifically to methods of wound healing using thymosin  $\beta$ 4.

**BACKGROUND OF THE INVENTION**

Inadequate methods and compositions to effectively heal chronic wounds  
15 is a significant health care problem. Impaired wound healing increases the chances of  
mortality and morbidity. This problem is especially prominent in patients with  
diabetes who develop severe, life threatening wounds on body extremities. Chronic  
diabetic foot ulcers often lead to amputations. These wounds are often the result of  
poor circulation derived from the diabetic patients' insulin-compromised cells as well  
20 as impaired vascularization of the wound bed, reduced infiltration of germ fighting  
cells and reduced tissue epithelialization. As a result, most current therapies include  
attempts to revascularize the wound bed and prevent infection.

Wounds in non-compromised tissues undergo a complex and ordered  
series of events to repair the tissue. The series of events may include infiltration of  
25 immune cells as part of the process to remove and destroy necrotic tissue, increased  
vascularization by angiogenic factors and increased cell proliferation and extracellular  
matrix deposition. Although the basic process of tissue repair has been characterized,

the individual steps and factors necessary to carry out this complex series of events are not well understood. The identification of individual steps and factors could lead to improved methods for the treatment of diseases resulting from inadequate wound repair processes.

5 Previous studies have used the "scratch" wound closure assay to assess the potential effects of an agent on *in vitro* cell migration. Though informative, such a test does not mimic the dynamic *in vivo* wound healing conditions to the extent that not all factors involved in wound closure are present in the *in vitro* assay. For this reason, *in vivo* systems have been developed to assess the ability of an agent or factor  
10 to modulate wound healing activities.

Using these types of *in vitro* models, a number of specific growth factors have been recognized for their effect on angiogenesis. One such growth factor is TGF- $\beta$ . This family of dimeric proteins includes TGF- $\beta$ 1, TGF- $\beta$ 2, TGF- $\beta$ 3, TGF- $\beta$ 4, and TGF- $\beta$ 5 which regulate the growth and differentiation of many cell types. This family of proteins exhibits a range of biological effects from stimulating the growth of some cell types (Noda *et al.*, (1989) *Endocrinology*, 124:2991-2995) and inhibiting the growth of other cell types (Goey *et al.*, (1989) *J. Immunol.*, 143:877-880; Pietenpol *et al.*, (1990) *Proc. Nat'l. Acad. Sci. USA*, 87:3758-3762). TGF- $\beta$  has also been shown to increase the expression of extracellular matrix proteins, including  
20 collagen and fibronectin (Ignatz *et al.*, (1986) *J. Biol. Chem.*, 261:4337-4345) and accelerates the healing of wounds (Mustoe *et al.*, (1987) *Science*, 237:1333-1335).

Another growth factor recognized for its effect on angiogenesis is Platelet Derived Growth Factor (PDGF). PDGF was originally found to be a potent mitogen for mesenchymal derived cells (Ross R. *et al.* (1974) *Proc Nat'l Acad Sci USA*  
25 71(4):1207-1210.; Kohler N. *et al.* (1974) *Exp. Cell Res.* 87:297-301). Further studies have shown that PDGF increases the rate of cellularity and granulation in tissue formation. Wounds treated with PDGF have the appearance of an early stage inflammatory response, including an increase in neutrophils and macrophage cell types at the wound site. These wounds also show enhanced fibroblast function  
30 (Pierce, GF *et al.* (1988) *J. Exp. Med.* 167:974-987). Both PDGF and TGF $\beta$  have

been shown to increase collagen formation, DNA content, and protein levels in animal studies. (Grotendorst, GR *et al.* (1985) *J. Clin. Invest.* 76:2323-2329.; Sporn, MB *et al.* (1983) *Science* 219:1329). The effect of PDGF in wound healing has been shown to be effective in human wounds. In human wounds, PDGF-AA expression is increased within pressure ulcers undergoing healing. The increase of PDGF-AA corresponds to an increase in activated fibroblasts, extracellular matrix deposition, and active vascularization of the wound. Furthermore, such an increase in PDGF-AA is not seen in chronic non-healing wounds. A number of other growth factors having the ability to induce angiogenesis and wound healing include, Vascular Endothelial Growth Factor (VEGF), Keratinocyte Growth Factor (KGF) and basic Fibroblast Growth Factor (bFGF).

However, most of these growth and angiogenic factors have side effects. Accordingly, there is a need for additional factors useful in promoting wound repair.

#### SUMMARY OF THE INVENTION

The present invention is based on the discovery that thymosin  $\beta 4$  (T $\beta 4$ ) accelerates wound healing and stimulates wound repair. Based on this finding, it is now possible to develop methods for accelerating wound healing in subjects having wounds in need of such treatment.

In a first embodiment, the invention provides a method for promoting wound repair in a subject in need of such treatment by administering to the subject or contacting the site of the wound with a wound-healing effective amount of a composition containing a wound healing polypeptide comprising the amino acid sequence LKKTET and conservative variants thereof having wound healing activity. In one aspect of the method, the wound healing polypeptide is T $\beta 4$  or an isoform of T $\beta 4$ .

In another embodiment, the invention provides a method for promoting tissue repair in a tissue in need of such treatment by contacting the tissue with an effective amount of a composition containing a wound healing polypeptide comprising the amino acid sequence LKKTET and conservative variants thereof

having wound healing activity, or nucleic acid encoding a wound healing polypeptide. In one aspect of the method, a wound healing peptide is T $\beta$ 4 or an isoform of T $\beta$ 4. The tissue may be contacted either *in vivo* or *ex vivo*.

In yet another embodiment, the invention provides a method of modulating wound repair in a subject in need of such treatment by systemic delivery of a wound-healing effective amount of a wound healing polypeptide comprising the amino acid sequence LKKTET and conservative variants thereof having wound healing activity. In one aspect of the method, a wound healing peptide is T $\beta$ 4 or an isoform of T $\beta$ 4.

In yet another embodiment, the present invention provides a method for stimulating epithelial cell migration at the site of a wound by contacting the wound with an effective amount of a T $\beta$ 4 polypeptide.

In another embodiment, the invention provides a method of diagnosing a pathological condition in a subject characterized by a wound healing disorder associated with T $\beta$ 4, including obtaining a sample suspected of containing T $\beta$ 4 from the subject, detecting a level of T $\beta$ 4 in the sample and comparing the level of T $\beta$ 4 with the level found in a normal sample (*i.e.*, a standard sample).

In another embodiment, the invention provides a method of ameliorating a wound healing disorder associated with T $\beta$ 4, including treating a subject having the disorder with a composition which modulates T $\beta$ 4 activity or the activity of a T $\beta$ 4 isoform.

In yet another embodiment, the present invention provides pharmaceutical compositions comprising a wound healing polypeptide comprising the amino acid sequence LKKTET and conservative variants thereof having wound healing activity and a pharmaceutically acceptable carrier. In one aspect, the wound healing polypeptide is T $\beta$ 4 or an isoform of T $\beta$ 4.

The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

## DESCRIPTION OF DRAWINGS

FIG. 1 is a schematic drawing of a wound.

FIG. 2 is a bar graph which shows the effect of topical and systemic delivery of T $\beta$ 4 on the width of a punch wound as compared to control. (A) Topical delivery of 5  $\mu$ g/50  $\mu$ l was performed on three of the six wounds in each animal on the day of wounding and at 48 hours after wounding. (B) Intraperitoneal injections of 60  $\mu$ g/300 $\mu$ l were done on the day of the wounding and thereafter every other day. Control animals were treated similarly with saline. Measurements are expressed as the mean percent decrease  $\pm$  SEM.

FIG. 3 is a bar graph which shows the effect of topical and systemic delivery of T $\beta$ 4 on the gap of a punch wound as compared to control. (A) Topical delivery of 5  $\mu$ g/50  $\mu$ l was performed on the day of wounding and at 48 hours after wounding. (B) Intraperitoneal injections of 60  $\mu$ g/300 $\mu$ l were done on the day of the wounding and thereafter every other day. Measurements are expressed as the mean percent decrease  $\pm$  SEM.

FIG. 4 is a histological section, stained with H&E, demonstrating the appearance of control and thymosin  $\beta$ 4 treated wounds at low magnification and higher magnification. Wounds are from day 7 as described in the legend to figure 2. Arrows indicate the edges of the original wound. (A) Control wound treated with saline. Migration of the epithelium is visible at the wound edges and debris are visible over the unhealed wound. (B) Increased re-epithelialization of the wound occurred when T $\beta$ 4 was injected intraperitoneally (60  $\mu$ g/300 $\mu$ l on alternate days). (C) Topical treatment (5 $\mu$ g/50 $\mu$ l of T $\beta$ 4) resulted in complete reepithelialization of the wound epidermis. Boxed areas are the location of the higher magnification fields (D-F). (D-F) Dermis near dermal and epidermal junction. (D) Control showing few cells near the dermis and little neovascularization. (E) and (F) Dermis showing granulation tissue infiltrated with fibroblasts and extensive neovascularization (arrowheads). (E) Intraperitoneal treatment and (F) topical application both resulted in significant new capillaries. (Scale bar = 1 mm).

FIG. 5 shows histological sections of 7 day wounds showing collagen deposition/accumulation. Masson's trichrome staining shows collagen and endothelial cells. (A) Low magnification view of a control wound treated with saline. (B) and (C). Low magnification views of wounds where T $\beta$ 4 was injected intraperitoneally (B) or applied topically (A). Boxed areas are the location of the higher magnification fields (D-F). Arrows indicate the edges of the original wound. (D) Control wound at higher magnification showing baseline collagen accumulation. Treatment intraperitoneally (E) or (F) topically resulted in enhanced collagen production/accumulation compared to wounds treated with saline. (Scale bar = 1 mm).

FIG. 6 shows T $\beta$ 4 stimulated keratinocyte migration in Boyden chamber assays. (A) T $\beta$ 4 in the lower wells of the chamber resulted in a 2-3 fold increase in migration on filters coated with collagen IV. The positive control, conditioned media, also showed increased migration over media alone.

FIG. 7 shows a graph demonstrating the migration of corneal epithelial cells at various concentrations of T $\beta$ 4.

FIG. 8 shows a graph representing corneal re-epithelialization in rat corneas in the presence and absence of T $\beta$ 4.

FIG. 9 shows a graph representing corneal re-epithelialization in the presence and absence of various concentrations of T $\beta$ 4.

FIG. 10 shows an amino acid sequence of T $\beta$ 4.

FIG. 11 shows the amino acid sequence of several known isoforms of T $\beta$ 4, and their phylogenetic distribution. N-terminal acetylation is indicated by "ac." Residues between 13 and 24 are thought to be important for actin binding.

## DETAILED DESCRIPTION OF THE INVENTION

Thymosin  $\beta$ 4 was initially identified as a protein that is up regulated during endothelial cell migration and differentiation *in vitro*. Thymosin  $\beta$ 4 was originally isolated from the thymus and is a 43 amino acid, 4.9 kDa ubiquitous polypeptide identified in a variety of tissues. Several roles have been ascribed to this

protein including a role in endothelial cell differentiation and migration, T cell differentiation, actin sequestration and vascularization. One biological activity of thymosin  $\beta 4$  (T $\beta 4$ ), as shown herein, effects tissue repair and wound healing. Another activity of T $\beta 4$  is anti-inflammatory activity.

5           The present invention resulted from investigation of the effects of T $\beta 4$  on wound healing. *In vivo* results have demonstrated that topical and systemic delivery of T $\beta 4$  promotes wound healing. Additional experiments demonstrated that T $\beta 4$ -treated wounds have increased extracellular matrix deposition in the wound bed.

10           The present invention identifies T $\beta 4$  as an active factor in promoting wound closure and tissue repair *in vivo* as well as increasing epithelial cell migration. *In vivo* administration of T $\beta 4$  indicates that cell migration, angiogenesis and extracellular matrix deposition are stimulated at or above the levels observed for migration, angiogenesis and matrix deposition in control animals. T $\beta 4$  promotes wound closure when administered systemically (*e.g.*, intra-peritoneally) and topically  
15 in wounded animal models. Increased levels of collagen were also observed in treated wounds showing that T $\beta 4$  treatment can also accelerate wound contraction and stimulate the healing process.

          The methods of the invention result from the identification of the effect of T $\beta 4$  on wound healing. *In vivo*, T $\beta 4$  stimulates wound healing in a full thickness  
20 punch wound (see Example 1) and in repair of eye-related wounds (Example 4). When given either topically or systemically (*e.g.*, intra-peritoneally) T $\beta 4$  accelerated closure and healing of wounds (see Example 1, 4, and 5).

#### **Promoting Tissue Regeneration**

25           In one embodiment, the invention provides a method for accelerating wound healing in a subject by contacting a wound with a wound-healing effective amount of a composition which contains T $\beta 4$  or a T $\beta 4$  isoform. The contacting may be topically or systemically. Examples of topical administration include, for example, contacting the wound with a lotion, salve, gel, cream, paste, spray, suspension, dispersion, hydrogel, ointment, or oil comprising T $\beta 4$ . Systemic administration

includes, for example, intravenous, intraperitoneal, intramuscular injections of a composition containing T $\beta$ 4 or a T $\beta$ 4 isoform. A subject may be any mammal, preferably human.

In addition, T $\beta$ 4 or a T $\beta$ 4 isoform is therapeutically valuable in cases where there is an impaired wound healing process, such as in wound healing compromised subjects. By "wound healing compromised" is meant subjects which have a reduced, decreased, or inability to recover from a wounding or trauma, due to recurrent wounding, trauma or inability of the subject's natural system to properly effectuate wound healing. For example, steroids reduce the ability of a subject to heal as compared to a subject which is not on steroids. Other such wounds present in compromised subjects include, but are not limited to, skin wounds such as diabetic ulcers, venus ulcers or pressure ulcers. Additionally, T $\beta$ 4 or a T $\beta$ 4 isoform is therapeutically valuable to augment the normal healing process.

As used herein, a "wound-healing effective amount" of a composition containing T $\beta$ 4 or a T $\beta$ 4 isoform for use in wound healing is defined as that amount that is effective in promoting tissue regeneration and repair. The "wound-healing effective amount" may be the therapeutically effective amount. Diseases, disorders or ailments possibly modulated by T $\beta$ 4 or a T $\beta$ 4 isoform include tissue repair subsequent to traumatic injuries or conditions including arthritis, osteoporosis and other musculo-skeletal disorders, burns, ulcers and other skin lesions, neurological and nerve disease and cardiovascular diseases including ischemia and atherosclerosis. Other potential tissues which can be treated by the methods and compositions of the invention include epidermal, eye, uro-genital, gastro-intestinal, cardiovascular, muscle, connective, and neural tissues. The term "induce", "induction" or "effect" as used herein, refers to the activation, stimulation, enhancement, initiation and/or maintenance of cellular mechanisms or processes necessary for the formation of a tissue or a portion thereof, repair process or tissue development as described herein.

### Modulation of Wound Healing

Wound healing, tissue regeneration and tissue repair result from a complex process that includes the proliferation and migration of inflammatory cells, endothelial cells, stromal cells and parenchymal cell, the deposition of extracellular matrix materials and the growth of new blood vessels, particularly capillaries. This complex process plays a crucial role in such beneficial functions as embryogenesis, the female reproductive cycle, as well as such abnormal functions as arthritis, chronic ulcerations and neuro-degenerative diseases.

In another embodiment, the invention provides a method for modulating wound healing in a subject or a tissue including contacting the subject or tissue with an effective wound-healing amount of a composition containing T $\beta$ 4 or a T $\beta$ 4 isoform. It is envisioned that T $\beta$ 4 or a T $\beta$ 4 isoform can be administered topically or systemically to prevent or treat a damaged tissue including, for example, tissues damaged due to ischemia, including ischemic brain, bone and heart disease, damage to corneal or retinal tissue of the eye, and damage to epithelial tissue, including skin.

In addition, the method of the invention is useful in promoting wound healing in tissues by promoting angiogenesis in tissue deprived of adequate blood flow. For example, a composition containing T $\beta$ 4 can promote the healing of chronic ulcers by increasing blood supply to the tissue site as well as increasing keratinocyte migration to close a wound.

T $\beta$ 4 isoforms have been identified and have about 70%, or about 75%, or about 80% or more homology to the amino acid sequence of T $\beta$ 4 set forth in Fig. 10. Such isoforms include, for example, T $\beta$ 4<sup>alt</sup>, T $\beta$ 9, T $\beta$ 10, T $\beta$ 11, T $\beta$ 12, T $\beta$ 13, T $\beta$ 14 and T $\beta$ 15 (Fig. 11; see also, Mihelic *et al.*, (1994) *Amino Acids*, 6:1-13, which describes the amino acid sequence of other T $\beta$ 4 isoforms, and is incorporated herein by reference). Similar to T $\beta$ 4, the T $\beta$ 10 and T $\beta$ 15 isoforms have been shown to sequester actin. T $\beta$ 4, T $\beta$ 10 and T $\beta$ 15, as well as these other isoforms share an amino acid sequence, LKKTET, that appears to be involved in mediating actin sequestration or binding. Although not wishing to be bound to any particular theory, the wound healing activity of T $\beta$ 4 and T $\beta$ 4 isoforms may be due, in part, to the ability to

polymerize actin. For example, T $\beta$ 4 can modulate actin polymerization in wounds to promote healing (*e.g.*,  $\beta$ -thymosins appear to depolymerize F-actin by sequestering free G-actin). T $\beta$ 4's ability to modulate actin polymerization may therefore be due to all, or in part, its ability to bind to or sequester actin via the LKKTET sequence.

- 5 Thus, as with T $\beta$ 4, other proteins which bind or sequester actin, or modulate actin polymerization, including T $\beta$ 4 isoforms having the amino acid sequence LKKTET, are likely to promote wound healing alone, or in a combination with T $\beta$ 4, as set forth herein.

10 Thus, it is specifically contemplated that known T $\beta$ 4 isoforms, such as T $\beta$ 4<sup>ala</sup>, T $\beta$ 9, T $\beta$ 10, T $\beta$ 11, T $\beta$ 12, T $\beta$ 13, T $\beta$ 14, and T $\beta$ 15, as well as T $\beta$ 4 isoforms not yet identified, will be useful in the methods of the invention. As such T $\beta$ 4 isoforms are useful in the methods of the invention, including the methods practiced in a subject, the invention therefore further provides pharmaceutical compositions comprising T $\beta$ 4 isoforms T $\beta$ 4<sup>ala</sup>, T $\beta$ 9, T $\beta$ 10, T $\beta$ 11, T $\beta$ 12, T $\beta$ 13, T $\beta$ 14, and T $\beta$ 15 and a pharmaceutically acceptable carrier.

15 In addition, other proteins having actin sequestering or binding capability, or that can mobilize actin or modulate actin polymerization, as demonstrated in an appropriate sequestering, binding, mobilization or polymerization assay, or identified by the presence of an amino acid sequence that mediates actin binding, such as LKKTET, for example, can similarly be employed in the methods of the invention. Such proteins include gelsolin, vitamin D binding protein (DBP), profilin, cofilin, depactin, DNaseI, vilin, fragmin, severin, capping protein,  $\beta$ -actinin and acumentin, for example. As such methods include those practiced in a subject, the invention further provides pharmaceutical compositions comprising gelsolin, vitamin D binding protein (DBP), profilin, cofilin, depactin, DNaseI, vilin, fragmin, severin, capping protein,  $\beta$ -actinin and acumentin as set forth herein. Thus, the invention includes the use of wound healing polypeptide comprising the amino acid sequence LKKTET and conservative variants thereof.

As used herein, the term "conservative variant" or grammatical variations thereof denotes the replacement of an amino acid residue by another, biologically similar residue. Examples of conservative variations include the replacement of a hydrophobic residue such as isoleucine, valine, leucine or methionine for another, the replacement of a polar residue for another, such as the substitution of arginine for lysine, glutamic for aspartic acids, or glutamine for asparagine, and the like.

T $\beta$ 4 has been localized to a number of tissue and cell types and thus, agents which stimulate the production of T $\beta$ 4 can be added to a composition to effect T $\beta$ 4 production from a tissue and/or a cell. Agents that effect wound repair can also be included in such a composition to augment the wound healing process. Such agents include members of the family of growth factors, such as insulin-like growth factor (IGF-1), platelet derived growth factor (PDGF), epidermal growth factor (EGF), transforming growth factor beta (TGF- $\beta$ ), basic fibroblast growth factor (bFGF), thymosin  $\alpha$ 1 (T $\alpha$ 1) and vascular endothelial growth factor (VEGF). More preferably, the agent is transforming growth factor beta (TGF- $\beta$ ) or other members of the TGF- $\beta$  superfamily. T $\beta$ 4 compositions of the invention aid in wound healing by effectuating growth of the connective tissue through extracellular matrix deposition, cellular migration and vascularization of the wound bed.

Additionally, agents that assist or stimulate the wound healing process may be added to a composition along with T $\beta$ 4 or a T $\beta$ 4 isoform to further modulate the wound healing process. Such agents include angiogenic agents, growth factors, agents that direct differentiation of cells, agents that promote migration of cells and agents that stimulate the provision of extracellular matrix materials in the wound bed. For example, and not by way of limitation, T $\beta$ 4 or a T $\beta$ 4 isoform alone or in combination can be added in combination with any one or more of the following agents: VEGF, KGF, FGF, PDGF, TGF $\beta$ , IGF-1, IGF-2, IL-1, prothymosin  $\alpha$  and thymosin  $\alpha$ 1 in a wound-healing effective amount.

In another aspect, the invention is useful for repair of tissue resulting from injuries due to surgical procedures, irradiation, laceration, toxic chemicals, viral infections, bacterial infections or burns. Additionally, the invention is useful for

revitalizing scar tissue resulting from any number of procedures, accidents or trauma. The term "scar tissue" means fibrotic or collagenous tissue formed during the healing of a wound or other morbid process. For example, T $\beta$ 4 can be included in a controlled release matrix which can be positioned in proximity to damaged tissue thereby promoting regeneration, repair and/or revascularization of such tissue. The term "controlled release matrix" means any composition that allows for the release of a bioactive substance which is mixed or admixed therein. The matrix can be a solid composition, a porous material (such as a scaffold, mesh, or sponge), or a semi-solid, gel or liquid suspension containing bioactive substances. The term "bioactive material" means any composition that modulates tissue repair when used in accordance with the method of the present invention. The bioactive materials or matrix can be introduced by means of injection, surgery, catheters or any other means suitable for modulating tissue repair.

It is envisioned that the methods and compositions of the invention can be used to aid wound healing and repair in guided tissue regeneration (GTR) procedures. Such procedures are currently used by those skilled in the medical arts to accelerate wound healing. Typically, nonresorbable or bioabsorbable materials are used to accelerate wound healing by promoting the repopulation of the wound area with cells which form the architectural and structural matrix of the tissue. For example, the methods and compositions of the invention can be used in aiding tissue repair or regeneration at an ulcer site in a human or other subject by placing a composition containing a bioresorbable polymer and T $\beta$ 4 at the site in need of tissue repair or regeneration such that the composition is effective for aiding tissue regeneration by releasing a wound-healing effective amount of T $\beta$ 4 at the site.

In another aspect, the invention is useful for the purposes of promoting tissue growth during the process of tissue engineering. As used herein, "tissue engineering" is defined as the creation, design, and fabrication of biological prosthetic devices, in combination with synthetic or natural materials, for the creation, augmentation or replacement of body tissues and organs. Thus, the present method can be used to augment the design and growth of human tissues outside the body, for

later implantation inside the body, or augment the design and growth of a tissue inside the body to repair or replace diseased or damaged tissue. For example, T $\beta$ 4 may be useful in promoting the growth of skin graft replacements which are used as a therapy in the treatment of burns and ulcers.

5 In another aspect of tissue engineering, T $\beta$ 4 can be included in external or internal devices containing human tissue designed to replace the function of a diseased internal tissue. This approach involves isolating cells from the body, placing them on or within a three-dimensional matrices and implanting the new system inside the body or using the system outside the body. The methods and compositions of the  
10 invention can be used and included in such matrices to promote the growth of tissues contained in the matrices. For example, T $\beta$ 4 can be included in a tissue engineered construct to promote the growth of the cells contained in the construct. It is envisioned that the method of the invention can be used to augment tissue repair, regeneration and engineering in endothelial cell-related products which may contain  
15 cartilage, cartilage-bone composites, bone, central nervous system tissues, muscle, liver, pancreatic islet (insulin-producing) cells, urogenital tissues, breast and tissues for gene therapy applications.

The present invention further provides methods and compositions for modulating female reproductive tract function. Growth factors have been shown to  
20 play a role in cyclic mitosis and differentiation of endometrial cellular components, recruitment of macrophages in decidualizing the endometrium, endometrial-trophoblast interactions, early pregnancy maintenance, and endometrial functional regeneration. The term "modulate" as used herein, denotes a modification of an existing condition or biologic state. Modulation of a condition as defined herein,  
25 encompasses both an increase or a decrease in the determinants affecting the existing condition. For example, administration of T $\beta$ 4 could be used to augment uterine functions in a condition where the promotion of endothelial cell growth is desired. For example, the uterus may be treated with T $\beta$ 4 to promote the growth and development of placental membranes or endometrial growth or the repair of these  
30 tissue following tissue injury. Furthermore, treatment with T $\beta$ 4 may be used to

promote and maintain a pregnancy by facilitating endometrial-trophoblast interaction. Alternatively, antagonist to T $\beta$ 4 could be administered to modulate conditions of excessive endometrial growth in which the level of T $\beta$ 4 is excessive in comparison to a normal biological condition. In addition, T $\beta$ 4 in combination with other agents, such as thymosin  $\alpha$ 1, may be desirable for treating disorders of the reproductive tract.

The therapeutic approaches described herein involve various routes of administration or delivery of reagents or compositions comprising the T $\beta$ 4 of the invention including any conventional administration techniques (for example, but not limited to, topical administration, local injection, inhalation, or systemic administration), to a subject with a wound or tissue in need of healing or repair.

Administration of T $\beta$ 4, as described above, can accelerate wound healing, increase cell migration into a wound site, induce the formation of tissue repair or regeneration, or promote the growth and development of the endometrium. The reagent, formulation or composition may also be targeted to specific cells or receptors by any method described herein or by any method known in the art of delivering, targeting T $\beta$ 4 polypeptides and expressing genes encoding T $\beta$ 4. For example, the methods and compositions using or containing T $\beta$ 4 of the invention may be formulated into pharmaceutical compositions by admixture with pharmaceutically acceptable non-toxic excipients or carriers. Such compositions may be prepared for parenteral administration, particularly in the form of liquid solutions or suspensions in aqueous physiological buffer solutions; for oral administration, particularly in the form of tablets or capsules; or for intranasal administration, particularly in the form of powders, nasal drops, or aerosols. Sustained release compositions are also encompassed by the present invention. Compositions for other routes of administration may be prepared as desired using standard methods.

A composition of the invention containing T $\beta$ 4 may be conveniently administered in unit dosage form, and may be prepared by any of the methods well known in the pharmaceutical art, for example, as described in Remington's Pharmaceutical Sciences (Mack Pub. Co., Easton, PA, 1990). Formulations for parenteral administration may contain as common excipients sterile water or saline,

polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes, and the like. In particular, biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxethylene-polyoxypropylene copolymers are examples of excipients for controlling the release of a compound of the invention *in vivo*. Other suitable parenteral delivery systems include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation administration may contain excipients such as lactose, if desired. Inhalation formulations may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or they may be oily solutions for administration in the form of nasal drops. If desired, the compounds can be formulated as a gel to be applied intranasally. Formulations for parenteral administration may also include glycocholate for buccal administration.

The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebrosides, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidyl-choline.

The targeting of liposomes has been classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific, and organelle-specific. Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries. Active

targeting, on the other hand, involves alteration of the liposome by coupling the liposome to a specific ligand such as a monoclonal antibody, sugar, glycolipid, or protein, or by changing the composition or size of the liposome in order to achieve targeting to organs and cell types other than the naturally occurring sites of localization.

The surface of the targeted delivery system may be modified in a variety of ways. In the case of a liposomal targeted delivery system, lipid groups can be incorporated into the lipid bilayer of the liposome in order to maintain the targeting ligand in stable association with the liposomal bilayer. Various linking groups can be used for joining the lipid chains to the targeting ligand. In general, the compounds bound to the surface of the targeted delivery system will be ligands and receptors which will allow the targeted delivery system to find and "home in" on the desired cells. A ligand may be any compound of interest which will bind to another compound, such as a receptor.

The therapeutic agents useful in the method of the invention can be administered parenterally by injection or by gradual perfusion over time. Administration may be intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, or transdermally.

Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents and inert gases and the like.

The invention also includes a pharmaceutical composition comprising a therapeutically effective amount of T $\beta$ 4 or a T $\beta$ 4 isoform in a pharmaceutically acceptable carrier. Such carriers include those listed above with reference to parenteral administration.

5           The actual dosage or reagent, formulation or composition that modulates a tissue repair process, fibrotic disorder, a sclerotic disorder, a cell proliferative disorder, or wound healing depends on many factors, including the size and health of a subject. However, one of ordinary skill in the art can use the following teachings describing the methods and techniques for determining clinical dosages (Spilker B.,

10 *Guide to Clinical Studies and Developing Protocols*, Raven Press Books, Ltd., New York, 1984, pp. 7-13, 54-60; Spilker B., *Guide to Clinical Trials*, Raven Press, Ltd., New York, 1991, pp. 93-101; Craig C., and R. Stitzel, eds., *Modern Pharmacology*, 2d ed., Little, Brown and Co., Boston, 1986, pp. 127-33; T. Speight, ed., *Avery's Drug Treatment: Principles and Practice of Clinical Pharmacology and Therapeutics*, 3d ed., Williams and Wilkins, Baltimore, 1987, pp. 50-56; R. Tallarida, R. Raffa and P. McGonigle, *Principles in General Pharmacology*, Springer-Verlag, New York, 1988, pp. 18-20) or to determine the appropriate dosage to use.

#### Antibodies that Bind to T $\beta$ 4

Antibodies to T $\beta$ 4 peptide or fragments could be valuable as diagnostic  
20 tools to aid in the detection of diseases in which T $\beta$ 4 is a pathological factor. Further, use of antibodies which bind to T $\beta$ 4 and inhibit or prevent the actions of T $\beta$ 4 are included in the present invention. Therapeutically, antibodies or fragments of the antibody molecule could also be used to neutralize the biological activity of T $\beta$ 4 in diseases where T $\beta$ 4 is over expressed. Such antibodies can recognize an epitope of  
25 T $\beta$ 4 or fragments thereof suitable for antibody recognition and neutralization of T $\beta$ 4 activity. As used in this invention, the term "epitope" refers to an antigenic determinant on an antigen, such as a T $\beta$ 4 peptide, to which the paratope of an antibody, such as an T $\beta$ 4-specific antibody, binds. Antigenic determinants usually consist of chemically active surface groupings of molecules, such as amino acids or

sugar side chains, and can have specific three-dimensional structural characteristics, as well as specific charge characteristics.

Preparation of an antibody requires a substantially purified moiety that can provide an antigenic determinant. The term "substantially pure" as used herein refers

to T $\beta$ 4, or variants thereof, which is substantially free of other proteins, lipids, carbohydrates or other materials with which it is naturally associated. Substantially purified or "isolated" refers to molecules, either nucleic or amino acid sequences, that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other components

with which they are naturally associated. One skilled in the art can isolate T $\beta$ 4 or a T $\beta$ 4 isoform using standard techniques for protein purification. The substantially pure peptide will yield a single major band on a non-reducing polyacrylamide gel. The purity of the T $\beta$ 4 peptide can also be determined by amino-terminal amino acid sequence analysis. T $\beta$ 4 or a T $\beta$ 4 isoform peptide includes functional fragments of the peptide, as long as the activity of T $\beta$ 4 or a T $\beta$ 4 isoform remains. Smaller peptides containing the biological activity of T $\beta$ 4 or a T $\beta$ 4 isoform are included in the invention. As used in the present invention, the term "antibody" includes, in addition to conventional antibodies, such protein fragments that have the ability to recognize specifically and bind the T $\beta$ 4 protein or variants thereof. Regions of the gene that

differ at the protein level are well defined. A protein can be raised by expression of the wild type (wt) gene or of the variants, or, preferably, fractions thereof. For example, the nucleic acid sequence can be cloned into expression vectors. According to this embodiment, the sequence of interest can first be obtained by employing PCR, as described above, or from a synthetic gene construction with overlapping and

ligated synthetic oligonucleotides. Another alternative would involve synthesis of a short peptide. All those methodologies are well known to one skilled in the art. See, for example, Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Volumes 1 and 2 (1987), with supplements, and Maniatis *et al.*, MOLECULAR CLONING, A LABORATORY MANUAL, Cold Spring Harbor

Laboratory, all of which are incorporated herein by reference.

The invention provides a method for detecting T $\beta$ 4, or variants thereof, which includes contacting an anti-T $\beta$ 4 antibody with a sample suspected of containing T $\beta$ 4, (e.g., cell or protein) and detecting binding to the antibody. An antibody which binds to T $\beta$ 4 peptide is labeled with a compound which allows  
5 detection of binding to T $\beta$ 4. There are many different labels and methods of labeling known to those of ordinary skill in the art. Examples of the types of labels which can be used in the present invention include enzymes, radioisotopes, fluorescent compounds, colloidal metals, chemiluminescent compounds, phosphorescent compounds, and bioluminescent compounds. Those of ordinary skill in the art will  
10 know of other suitable labels for binding to the antibody, or will be able to ascertain such, using routine experimentation. For purposes of the invention, an antibody specific for T $\beta$ 4 peptide may be used to detect the level of T $\beta$ 4 in biological fluids and tissues. Any specimen containing a detectable amount of antigen can be used. The level of T $\beta$ 4 in the suspect cell can be compared with the level in a normal cell to determine whether the subject is predisposed to a T $\beta$ 4 associated increase in angiogenesis or wound healing.

Use of antibodies for the diagnostic methods of the invention includes, for example, immunoassays in which the antibodies can be utilized in liquid phase or bound to a solid phase carrier. In addition, the antibodies in these immunoassays can  
20 be detectably labeled in various ways. Examples of types of immunoassays which can utilize antibodies of the invention are competitive and non-competitive immunoassays in either a direct or indirect format. Examples of such immunoassays are the radioimmunoassay (RIA) and the sandwich (immunometric) assay. Detection of the antigens using the antibodies of the invention can be done utilizing  
25 immunoassays which are run in either the forward, reverse, or simultaneous modes, including immunohistochemical assays on physiological samples. Those of skill in the art will know, or can readily discern, other immunoassay formats without undue experimentation.

T $\beta$ 4 antibodies can be bound to many different carriers and used to detect the presence of an antigen comprising the peptide of the invention. Examples of well-known carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, agaroses and magnetite. The nature of the carrier can be either soluble or insoluble for purposes of the invention. Those skilled in the art will know of other suitable carriers for binding antibodies, or will be able to ascertain such, using routine experimentation.

Another technique which may also result in greater sensitivity consists of coupling the antibodies to low molecular weight haptens. These haptens can then be specifically detected by means of a second reaction. For example, it is common to use such haptens as biotin, which reacts with avidin, or dinitrophenyl, puridoxal, and fluorescein, which can react with specific antihapten antibodies.

The invention includes use of antibodies immunoreactive with T $\beta$ 4 peptide or functional fragments thereof. Antibody which consists essentially of pooled monoclonal antibodies with different epitopic specificities, as well as distinct monoclonal antibody preparations are provided. Monoclonal antibodies are made from antigen containing fragments of the protein by methods well known to those skilled in the art (Kohler, *et al.*, *Nature*, 256:495, 1975). The term antibody as used in this invention is meant to include intact molecules as well as fragments thereof, such as Fab and F(ab')<sub>2</sub>, Fv and SCA fragments which are capable of binding an epitopic determinant on T $\beta$ 4.

(1) An Fab fragment consists of a monovalent antigen-binding fragment of an antibody molecule, and can be produced by digestion of a whole antibody molecule with the enzyme papain, to yield a fragment consisting of an intact light chain and a portion of a heavy chain.

(2) An Fab' fragment of an antibody molecule can be obtained by treating a whole antibody molecule with pepsin, followed by reduction, to yield a molecule consisting of an intact light chain and a portion of a heavy chain. Two Fab' fragments are obtained per antibody molecule treated in this manner.

(3) An (Fab')<sub>2</sub> fragment of an antibody can be obtained by treating a whole antibody molecule with the enzyme pepsin, without subsequent reduction. A (Fab')<sub>2</sub> fragment is a dimer of two Fab' fragments, held together by two disulfide bonds.

(4) An Fv fragment is defined as a genetically engineered fragment containing the variable region of a light chain and the variable region of a heavy chain expressed as two chains.

(5) A single chain antibody ("SCA") is a genetically engineered single chain molecule containing the variable region of a light chain and the variable region of a heavy chain, linked by a suitable, flexible polypeptide linker.

Alternatively, a therapeutically or diagnostically useful anti-Tß4 antibody may be derived from a "humanized" monoclonal antibody. Humanized monoclonal antibodies are produced by transferring mouse complementary determining regions from heavy and light variable chains of the mouse immunoglobulin into a human variable domain, and then substituting human residues in the framework regions of the murine counterparts. The use of antibody components derived from humanized monoclonal antibodies obviates potential problems associated with the immunogenicity of murine constant regions. General techniques for cloning murine immunoglobulin variable domains are described, for example, by Orlandi *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 3833 (1989), which is hereby incorporated in its entirety by reference. Techniques for producing humanized monoclonal antibodies are described, for example, by Jones *et al.*, *Nature* 321: 522 (1986); Riechmann *et al.*, *Nature* 332: 323 (1988); Verhoeven *et al.*, *Science* 239: 1534 (1988); Carter *et al.*, *Proc. Nat'l Acad. Sci. USA* 89: 4285 (1992); Sandhu, *Crit. Rev. Biotech.* 12: 437 (1992); and Singer *et al.*, *J. Immunol.* 150: 2844 (1993), which are hereby incorporated by reference.

Antibodies of the invention also may be derived from human antibody fragments isolated from a combinatorial immunoglobulin library. See, for example, Barbas *et al.*, *METHODS: A COMPANION TO METHODS IN ENZYMOLOGY*, VOL. 2, page 119 (1991); Winter *et al.*, *Ann. Rev. Immunol.* 12: 433 (1994), which are hereby incorporated by reference. Cloning and expression vectors that are useful

for producing a human immunoglobulin phage library can be obtained, for example, from STRATAGENE Cloning Systems (La Jolla, CA).

**Methods and Compositions for Treating or Diagnosing T $\beta$ 4-Associated Disorders**

In another embodiment of the invention, a method of diagnosing a  
5 pathological state in a subject suspected of having a pathology characterized by a disorder associated with T $\beta$ 4 is provided. The method includes obtaining a sample suspected of containing T $\beta$ 4 from the subject, determining the level of T $\beta$ 4 in the sample and comparing the level of T $\beta$ 4 in the sample to the level of T $\beta$ 4 in a normal standard sample. Such conditions include, but are not limited to subjects having cell  
10 proliferative disorders, recurrent wounds, tissue repair disorders, fibrotic tissue disorders, chronic ulcers and other disorders described herein. Such disorders further include those associated with the various T $\beta$ 4 isoforms, known or not yet identified.

The term "cell-proliferative disorder" denotes malignant as well as non-malignant cell populations which often appear to differ from the surrounding tissue  
15 both morphologically and genotypically. Malignant cells (*i.e.* cancer) develop as a result of a multistep process. Such disorders may be detected using the methods of the current invention. For example, a sample suspected of containing T $\beta$ 4 is obtained from a subject, the level of T $\beta$ 4 peptide is determined and compared with the level of T $\beta$ 4 peptide in a normal tissue sample. The level of T $\beta$ 4 can be determined by any  
20 number of methods including, for example, immunoassay using anti-T $\beta$ 4 peptide antibodies. Other variations of such assays include radioimmunoassay (RIA), ELISA and immunofluorescence. Alternatively, nucleic acid probes can be used to detect and quantify T $\beta$ 4 peptide mRNA for the same purpose. Such detection methods are standard in the art.

25 In another embodiment, the invention provides a method for ameliorating a wound healing disorder associated with T $\beta$ 4 or a T $\beta$ 4 isoform, including treating a subject having the disorder with a composition that regulates T $\beta$ 4 activity. The term "ameliorate" denotes a lessening of the detrimental effect of the disease-inducing response in the subject receiving therapy. Where the disease is due to an abnormally

high level of T $\beta$ 4, the administration of an agent, such as an antagonist of T $\beta$ 4 activity, may be effective in decreasing the amount of T $\beta$ 4 activity. Alternatively, where the disease is due to an abnormally low level of T $\beta$ 4, the administration of T $\beta$ 4 or an agent that increases T $\beta$ 4 activity, such as an agonist, may be effective in  
5 increasing the amount of T $\beta$ 4 activity.

In yet another embodiment, the invention provides a method of treating a subject having a wound healing disorder characterized by recurrent or slow to heal wounds or wounds that are chronic non-healing wounds associated with altered T $\beta$ 4 or T $\beta$ 4 isoform gene expression in a subject. The method includes administering to a  
10 subject having the disorder a wound-healing effective amount of an agent which modulates T $\beta$ 4 gene expression, thereby treating the disorder. The term "modulate" refers to inhibition or suppression of T $\beta$ 4 expression when T $\beta$ 4 is over expressed, and induction of expression when T $\beta$ 4 is under expressed. The term "wound-healing effective amount" means that amount of T $\beta$ 4 agent which is effective in modulating  
15 T $\beta$ 4 gene expression resulting in reducing the symptoms of the T $\beta$ 4 associated wound healing disorder.

An agent which modulates T $\beta$ 4 or T $\beta$ 4 isoform gene expression may be a polynucleotide for example. The polynucleotide may be an antisense, a triplex agent, or a ribozyme. For example, an antisense may be directed to the structural gene  
20 region or to the promoter region of T $\beta$ 4 may be utilized.

When a wound healing disorder is associated with the expression of T $\beta$ 4, a therapeutic approach which directly interferes with the translation of T $\beta$ 4 mRNA into protein is possible. For example, an antisense nucleic acid or a ribozyme can be used to bind to the T $\beta$ 4 RNA or to cleave it. Antisense RNA or DNA molecules bind  
25 specifically with a targeted gene's RNA message, interrupting the expression of that gene's protein product. The antisense binds to the mRNA forming a double stranded molecule which cannot be translated by the cell. Antisense oligonucleotides of about 15-25 nucleotides are preferred since they are easily synthesized and have an inhibitory effect just like antisense RNA molecules. In addition, chemically reactive  
30 group, such as iron-linked ethylenediaminetetraacetic acid (EDTA-Fe) can be

attached to an antisense oligonucleotide, causing cleavage of the RNA at the site of hybridization. These and other uses of antisense methods to inhibit the *in vitro* translation of genes are well known in the art (Marcus-Sakura, *Anal., Biochem.*, 172:289, 1988).

Antisense nucleic acids are DNA or RNA molecules that are complementary to at least a portion of a specific mRNA molecule (Weintraub, *Scientific American*, 262:40, 1990). In the cell, the antisense nucleic acids hybridize to the corresponding mRNA, forming a double-stranded molecule. The antisense nucleic acids interfere with the translation of the mRNA, since the cell will not translate a mRNA that is double-stranded. Antisense oligomers of about 15 nucleotides are preferred, since they are easily synthesized and are less likely to cause problems than larger molecules when introduced into the target T $\beta$ 4 producing cell. The use of antisense methods to inhibit the *in vitro* translation of genes is well known in the art (Marcus-Sakura, *Anal. Biochem.*, 172:289, 1988).

Use of an oligonucleotide to stall transcription is known as the triplex strategy since the oligomer winds around double-helical DNA, forming a three-strand helix. Therefore, these triplex compounds can be designed to recognize a unique site on a chosen gene (Maher, *et al.*, *Antisense Res. and Dev.*, 1(3):227, 1991; Helene, C., *Anticancer Drug Design*, 6(6):569, 1991).

Ribozymes are RNA molecules possessing the ability to specifically cleave other single-stranded RNA in a manner analogous to DNA restriction endonucleases. Through the modification of nucleotide sequences which encode these RNAs, it is possible to engineer molecules that recognize specific nucleotide sequences in an RNA molecule and cleave it (Cech, *J. Amer. Med. Assn.*, 260:3030, 1988). A major advantage of this approach is that, because they are sequence-specific, only mRNAs with particular sequences are inactivated.

There are two basic types of ribozymes namely, *tetrahymena*-type (Hasselhoff, *Nature*, 334:585, 1988) and "hammerhead"-type. *Tetrahymena*-type ribozymes recognize sequences which are four bases in length, while "hammerhead"-type ribozymes recognize base sequences 11-18 bases in length. The longer the

recognition sequence, the greater the likelihood that the sequence will occur exclusively in the target mRNA species.

These and other uses of antisense methods to inhibit the *in vivo* translation of genes are well known in the art (e.g., De Mesmaeker, *et al.*, 1995. Backbone

- 5 modifications in oligonucleotides and peptide nucleic acid systems. *Curr. Opin. Struct. Biol.* 5:343-355; Gewirtz, A.M., *et al.*, 1996b. Facilitating delivery of antisense oligodeoxynucleotides: Helping antisense deliver on its promise; *Proc. Natl. Acad. Sci. U.S.A.* 93:3161-3163; Stein, C.A. A discussion of G-tetrads 1996. Exploiting the potential of antisense: beyond phosphorothioate
- 10 oligodeoxynucleotides. *Chem. and Biol.* 3:319-323).

- Delivery of antisense, triplex agents, ribozymes, competitive inhibitors and the like can be achieved using a recombinant expression vector such as a chimeric virus or a colloidal dispersion system. Various viral vectors which can be utilized for gene therapy as taught herein include adenovirus, herpes virus, vaccinia, or, preferably, an RNA virus such as a retrovirus. Preferably, the retroviral vector is a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). A number of
- 20 additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a polynucleotide sequence of interest into the viral vector, along with another gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is now target specific. Retroviral vectors
- 25 can be made target specific by inserting, for example, a polynucleotide encoding a sugar, a glycolipid, or a protein. Preferred targeting is accomplished by using an antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of
- 30 the retroviral vector containing the antisense polynucleotide.

Since recombinant retroviruses are defective, they require assistance in order to produce infectious vector particles. This assistance can be provided, for example, by using helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR.

- 5 These plasmids are missing a nucleotide sequence which enables the packaging mechanism to recognize an RNA transcript for encapsidation. Helper cell lines which have deletions of the packaging signal include but are not limited to  $\Psi$ 2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging
- 10 signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced.

Alternatively, NIH 3T3 or other tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes *gag*, *pol* and *env*, by conventional calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The resulting cells release the retroviral vector into the culture medium.

- A targeted delivery system for delivery of nucleic acids as described herein includes a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, gene activated
- 20 matrices and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles *in vitro* and *in vivo*. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0  $\mu$ m can encapsulate a substantial percentage of an
- 25 aqueous buffer containing large macromolecules. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, *et al.*, *Trends Biochem. Sci.*, 6:77, 1981). In addition to mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells. In order for a liposome to be an efficient gene transfer
- 30 vehicle, the following characteristics should be present: (1) encapsulation of the genes

of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information

5 (Mannino, *et al.*, *Biotechniques*, 6:682, 1988).

Pathologically, T $\beta$ 4 may be involved in diseases in which there is an overgrowth of blood vessels, such as cancer, tumor formation and growth, diabetic retinopathy, neovascular glaucoma, rheumatoid arthritis and psoriasis.

10 The ingrowth of capillaries and ancillary blood vessels is essential for growth of solid tumors and is thus an unwanted physiological response which facilitates the spread of malignant tissue and metastases. Inhibition of angiogenesis and the resultant growth of capillaries and blood vessels is therefore a component of effective treatment of malignancy in use of treatment of cancer patients.

Thus, in another embodiment, the invention provides a method of inhibiting angiogenesis in a subject, including administering to the subject a composition containing an agent which regulates T $\beta$ 4 activity. The composition may include agents that regulate angiogenesis, for example agents that affect thymosin  $\alpha$ 1, PDGF, VEGF, IGF, FGF and TGF $\beta$ . For example, the inhibition of angiogenesis and endothelial cell migration can be beneficial in controlling the growth of solid tumors.

20 Most, if not all solid tumors, like normal tissue, require a steady and sufficient blood supply for optimal growth. Tumors are known to make use of angiogenic growth factors to attract new blood vessels and ascertain supply with sufficient amounts of nutrients to sustain their growth. Many tumors are well vascularized and the inhibition of the formation of an adequate blood supply to the tumor by inhibition of tumor vascularization, as a result of inhibition of angiogenesis, is beneficial in tumor growth control. Without a strong blood supply, rapid and prolonged growth of tumor tissue cannot be sustained. Thus, agents that inhibit T $\beta$ 4 activity may be used to prevent neoplastic growth. The T $\beta$ 4 inhibiting agent may be administered orally, parenterally, topically, intravenously, or systemically. In addition, for inhibiting

30 tumor cell proliferation and tumor growth, the agent may be administered locally

directly to the tumor or as a part of a deposited slow release formulation.

Administration may be on a daily basis for as long as needed to inhibit angiogenesis, endothelial cell proliferation, tumor cell proliferation or tumor growth. Alternatively, a slow release formulation may continue for as long as needed to control tumor

5 growth. This dosage regimen may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

In this regard, the compositions of this invention that are useful as  
10 inhibitors of angiogenesis, endothelial cell proliferation, tumor cell proliferation and tumor growth contain a pharmaceutically acceptable carrier and an amount of T $\beta$ 4 modulating agent effective to inhibit tumor or endothelial cell proliferation. Such compositions may also include preservatives, antioxidants, immunosuppressants and other biologically and pharmaceutically effective agents which do have effects on tumor growth but which do not exert a detrimental effect on the T $\beta$ 4 modulating agent. For treatment of tumor cells the composition may include a chemotherapeutic agent, for example an anti-cancer agent which selectively kills the faster replicating tumor cells, many of which are known and clinically used. Exemplary anticancer agents include mephalan, cyclophosphamide, methotrexate, adriamycin and  
20 bleomycin.

#### Screen for compounds which modulate T $\beta$ 4 activity

In another embodiment, the invention provides a method for identifying a compound that modulates T $\beta$ 4 activity, angiogenesis activity or wound healing activity. The method includes incubating components including the compound and  
25 T $\beta$ 4 under conditions sufficient to allow the components to interact and determining the effect of the compound on T $\beta$ 4 activity before and after incubating in the presence of the compound. Compounds that affect T $\beta$ 4 activity (e.g., antagonists and agonists) include peptides, peptidomimetics, polypeptides, chemical compounds, minerals such

as zincs, and biological agents. T $\beta$ 4 activity can be assayed using the methodology as described in the present Examples.

The present Examples are meant to illustrate, but not limit the scope of the appended claims. Accordingly, one skilled in the art will recognize a number of equivalent materials and methods, which are intend to be covered by the present invention and disclosure.

### EXAMPLE 1

#### In vivo wound healing is accelerated by T $\beta$ 4

T $\beta$ 4, whether administered topically or intraperitoneal, significantly accelerated wound healing as compared to untreated wounds (Fig. 2 and 3). Full thickness 8 mm punch biopsy wounds were made on the dorsal surface of rats as previously reported (Bhartiya *et al.*, *J. Cell. Physiol.* 150:312, 1992; Sihhu *et al.*, *J. Cell. Physiol.* 169:108, 1996) and T $\beta$ 4 was given topically at the time of wounding (5  $\mu$ g in 50  $\mu$ l) and again after 48 hours. Controls for the topical treatment received identical amounts of saline at the time of wounding and at 48 hours. Additional rats received intraperitoneal injections at the time of wounding (60  $\mu$ g in 300  $\mu$ l) and again every other day (*e.g.*, days 0, 2, 4, and 6). Controls for these animals received identical amounts of saline intra-peritoneally on the same injection schedule. On days 4 and 7 post-wounding, measurements were made on the wound size. At days 8 and 9 post-wounding, tissue was collected and fixed in 10% buffered formalin. The samples were sectioned and stained with H&E and Masson's Trichrome (American Histolabs, Gaithersburg, MD).

Histological sections were used to measure the re-epithelialization and the contraction of the wound using an ocular micrometer. Epidermal migration was determined by measuring the lengths of the tongues of epithelium migrating from either side of the wound over the wound bed from the zone of proliferation at the margin of the uninjured and wounded skin. Epidermal thickness was also measured beginning at the junction of the uninjured and proliferating epidermis. The thickness was measured vertically from the basement membrane to the most superficial layer of

the migrating epidermis at every 200 microns. The mean epidermal thickness of each migrating tongue of epidermis was then computed from each wound. Vessel counts were performed by first identifying vascular spaces by their endothelial lining. All such vessels in the wound bed were counted including those at the junction of the dermis and the subcutis, since angiogenesis into the wounds occurs to a great extent from these vessels. The numbers were averaged into vessel counts per 10 high powered fields (40x).

The effect of T $\beta$ 4 on wound healing was studied in a full thickness cutaneous rat wound model. FIG. 1 shows a diagram of the wound site that extends from the epidermis to the fat/muscle layer. This model allowed measurement of two parameters: the re-epithelialization (gap) and the contraction (width) of the wound. Wounds treated topically with T $\beta$ 4 showed about a 15% decrease in width and about 15% decrease in gap in the treated versus controls (FIG. 2 and 3, respectively).

Figure 2 shows a 15% decrease in wound width as compared to the saline controls as early as 4 days after wounding and continued until day 7. Intraperitoneal injection of T $\beta$ 4 resulted in a 18% decrease in wound width relative to saline treated controls at day 4 and 11% decrease at day 7. This trend was observed on the 4th day post wounding and continued through day 7 (\* $P \leq 0.0001$ , \*\* $P \leq 0.08$ , significant difference from media alone, student's t-test). These data demonstrate that T $\beta$ 4, when given either topically or systemically, increases wound re-epithelialization and contraction. Both topical and systemic treatment are equally effective. Lower doses of T $\beta$ 4 were tested including 2.5  $\mu$ g and 0.5  $\mu$ g in 50  $\mu$ l for topical and 30  $\mu$ g and 6  $\mu$ g in 300  $\mu$ l for intraperitoneal injection but reduced or no effect, respectively, was observed on wound healing.

Figure 3 shows an 18% decrease in gap length as compared to saline controls when T $\beta$ 4 is administered topically, as early as 4 days after wounding. This trend continued to termination at day 7 (\* $P \leq 0.04$ , student's t-test). Intraperitoneal injections resulted in a 42% decrease in gap size relative to saline treated controls. This decrease was observed on the 4th day post wounding and continued through day 7 (\*\* $P \leq 0.0007$ , student's t-test). The increase in re-epithelialization was observed in

wounds treated for 7 days and the rate of gap closure was slightly accelerated over that observed at day 4. A 62% decrease in gap size was observed in the T $\beta$ 4-treated wounds. Quantitation of epidermal migration showed a statistically significant 1.5 fold increase in migration of epidermal tongues over the wound bed after topical treatment (Table 1). Quantitation of epithelial migration in intraperitoneally treated wounds showed a statistically significant increase in migration of epidermal tongues as compared to controls (Table 1). There was no difference in the thickness of the migrating epidermis between either of the T $\beta$ 4 treatments and the control (Table 1). Histological sections of the wounds clearly show increased re-epithelialization in the treated wounds as compared to controls in 7 day wounds (FIG 4).

Table 1: Morphometric Measurements of Control and Thymosin  $\beta$ 4 Treated Samples

Parameter	Control	I.P.	Topical
Epidermal Migration ( $\mu$ m)	2403.3 $\pm$ 9.7	3168.3 $\pm$ 38.4*	3668.7 $\pm$ 56.6*
Epidermal Thickness ( $\mu$ m)	128.2 $\pm$ 19.3	135.0 $\pm$ 11.7	142.3 $\pm$ 19.8
Vessels/10 HPF	1364.0 $\pm$ 15.0	2415.0 $\pm$ 24.3*	2186.0 $\pm$ 11.8*

HPF: high power field. \*P $\leq$ 0.00001 by Welch's t-test, significantly different than control.

FIG. 4 shows a comparison of typical control (D) and T $\beta$ 4-treated (E and F) sections of 7 day wounds. Treatment with T $\beta$ 4 resulted in considerable capillary ingrowth (FIG 4E and F, arrows). Vessel counts showed a significant (about 2 fold) increase in the number of vessels in T $\beta$ 4 treated wounds (Table 1). No increases in the number of macrophages in the wounds were observed. There was no apparent increase in the accumulation/biosynthesis of collagen in treated -T $\beta$ 4 wounds (Fig. 5B and C vs A) supporting a decreased wound width and supporting a role for T $\beta$ 4 in wound contraction. Both the topical and systemically treated wound appeared similar although the wound contraction proceeded slightly more quickly with the topical treatment.

Reduction of the wound size was observed in both experimental groups as compared to control groups (Fig. 2 - 4). More and larger blood vessels were noted in the experimental groups as compared to the controls (Fig. 4). Additionally, an increase in the accumulation/biosynthesis of collagen by T $\beta$ 4 treated wounds as compared to control suggests a role for T $\beta$ 4 in wound contraction and extracellular matrix deposition. Histological staining of these wounds demonstrated an increase in collagen density and extracellular matrix deposition when compared to controls. (Fig. 5).

## EXAMPLE 2

### Migration Assays of Keratinocytes

Primary keratinocytes were prepared from either Balb/c or CD-1 newborn mice as described previously (Dlugosz *et al.*, 1995). Cells were plated in calcium- and magnesium-free Eagle's Minimal Essential Medium (EMEM) containing 8% fetal calf serum treated with 8% Chelex (Bio-Rad Laboratories, Hercules, CA), 20 units/ml penicillin-streptomycin, and the calcium concentration was adjusted to 0.25 mM. The following day, cultures were washed with calcium- and magnesium-free phosphate buffered saline, treated briefly with Trypsin (Life Technologies, Gaithersburg, MD), washed with culture medium and resuspended in EMEM containing 0.05 mM calcium. Cells were used immediately in migration assays.

Keratinocyte migration assays were carried out in Boyden chamber using 12  $\mu$ m pore polyester membranes (Poretics, Livermore, CA) coated with a 0.1 mg/ml solution of collagen IV in dH<sub>2</sub>O (Trevigen, Gaithersburg, MD). Filters were then dried at least 1 h. Cells were harvested using Versene or Trypsin (Life Technologies, Gaithersburg, MD) and resuspended in Eagle's minimal essential medium with 0.05 mM Ca<sup>2+</sup>. The bottom chamber was loaded with EMEM containing 0.01, 0.1, 10, 100, and 1000 ng/ml of synthetic T $\beta$ 4. Conditioned medium from primary dermal fibroblasts and/or keratinocyte growth factor was added to several wells as a positive control. Cells were added to the upper chamber at a concentration of 50,000 cells per well. Chambers were incubated at 35 C/7% CO<sub>2</sub> for 4-5 hours and the filters were

then fixed and stained using Diff-Quik (Baxter Healthcare Corporation, McGraw Park, IL). The cells that migrated through the filter were quantitated by counting the center of each well at 10x using an Olympus CK2 microscope. Each condition was assayed in triplicate wells and each experiment was repeated four times with different preparations of cells.

The results demonstrated that keratinocyte migrated in response to T $\beta$ 4 after 4-5 hours of exposure. Migration was enhanced 2-3 fold ( $P \leq 0.003$ ) over migration in the presence of media alone (FIG. 6) and at the maximal responding dose exceeded the positive control. The effect of T $\beta$ 4 on migration, while showing slightly different dose curves depending on the cell preparation and source, clearly showed a biphasic pattern with 1000 ng/ml and 0.01 ng/ml showing the most migration and the middle doses showing less stimulation (but still greater than control media) in all 4 assays.

### EXAMPLE 3

#### Migration Assays of Corneal Epithelial Cells

Corneal Epithelial Cell migration assays were carried out in Boyden chamber using 12  $\mu$ m pore polyester membranes (Poretics, Livermore, CA) coated with a 0.1 mg/ml solution of collagen IV in dH<sub>2</sub>O (Trevigen, Gaithersburg, MD). Filters were then dried at least 1 h. Cells were cultured and resuspended in Eagle's Minimal Essential Medium with 0.05 mM Ca<sup>2+</sup>. The bottom chamber was loaded with EMEM containing 0.01, 0.1, 10, 100, and 1000 ng/ml of synthetic T $\beta$ 4. Conditioned medium from primary dermal fibroblasts and/or keratinocyte growth factor was added to several wells as a positive control. Cells were added to the upper chamber at a concentration of 50,000 cells per well. Chambers were incubated at 35 C/7% CO<sub>2</sub> for 4-5 hours and the filters were then fixed and stained using Diff-Quik (Baxter Healthcare Corporation, McGraw Park, IL). The cells that migrated through the filter were quantitated by counting the center of each well at 10x using an Olympus CK2 microscope. Each condition was assayed in triplicate wells and each experiment was repeated four times with different preparations of cells.

The results demonstrated that corneal epithelial cell migrated in response to T $\beta$ 4 after 4-5 hours of exposure. Migration was enhanced 2-3 fold over migration in the presence of media alone (FIG. 7) with the highest level of migration seen at 100 ng/ml of T $\beta$ 4.

#### EXAMPLE 4

##### In vivo Corneal Re-Epithelialization

To determine the effect of T $\beta$ 4 on corneal re-epithelialization *in vivo*, Rat corneas were de-epithelialized and treated with T $\beta$ 4. Filters were soaked in heptanol, applied to the eye for 30 seconds, and then the epithelium was scraped. Various concentration of T $\beta$ 4 in saline was applied to the eye and at 24 hours the rats were sacrificed. The eyes were fixed, sectioned and the degree of corneal epithelial migration (as measured in pixels) was determined using a microscope with an internal caliper by a masked observer. The results demonstrate that re-epithelialization of the cornea was increased 2-fold over untreated control in the presence of about 1 to 25 $\mu$ g of T $\beta$ 4 (FIG. 8 and 9). In addition, it was noted that T $\beta$ 4 treated eyes had reduced inflammation compared to the non-treated corneas.

#### EXAMPLE 5

##### Impaired Healing Model

Thymosin  $\beta$ 4 also enhanced wound healing in an impaired model. Steroid treatment reduces the rate of wound repair in mammals. Rats treated with steroids such as hydrocortisone serve as a model of impaired wound healing due to the delay observed in wound closure. Animals were injected intramuscularly everyday with hydrocortisone. Steroid treated rats showed a significant increase in the level of healing when T $\beta$ 4 was added topically or injected intraperitoneally. At the initial time point, day 4, topically treated animals showed little response ( $\leq 7\%$  gap or width closure, N=3) compared to saline treatment. Intraperitoneal injection, however, resulted in a 28% decrease in gap size and a 14% decrease in wound width. At day 7, a response was observed with both topical treatment and intraperitoneal injection.

The gap in topically treated animals decreased by 39% compared to saline treatment. The wound width decreased by 23%. Intraperitoneal injection resulted in a 26% decrease in gap size and a 10% decrease in wound width. Taken together, these demonstrate that T $\beta$ 4 is useful to treat chronic, as well as, acute wounds.

- 5        A number of embodiments of the present invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention. Accordingly, other embodiments are within the scope of the following claims.

0972445-01901